

**Doptelet**<sup>®</sup>  
(avatrombopag)  
tablets

**Doptelet**<sup>®</sup>  
**Sprinkle**  
(avatrombopag) oral granules

STEP RIGHT UP TO LEARN ABOUT  
AN EXPANDED INDICATION



FOLLOW ME, I'M HERE  
TO HELP YOU LEARN MORE!

## HERE'S WHAT YOU SHOULD KNOW BEFORE YOUR CHILD STARTS DOPTELET...

### What is DOPTELET and DOPTELET Sprinkle (avatrombopag)?

Doptelet is a prescription medicine used to treat low blood platelet counts in:

- children 1 year and older with persistent or chronic ITP when other treatments have not worked well enough

DOPTELET and DOPTELET Sprinkle are different dosage forms of DOPTELET.

DOPTELET and DOPTELET Sprinkle are not used to make platelet counts normal.

It is not known if DOPTELET or DOPTELET Sprinkle are safe and effective in children younger than 1 year of age.

### IMPORTANT SAFETY INFORMATION

**Before you take DOPTELET or DOPTELET Sprinkle**, tell your doctor about all your medical conditions, including if you:

- Have ever had a blood clot

**Tell your healthcare provider about all of the medicines you take, including** prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTELET or DOPTELET Sprinkle may affect the way other medicines work, and other medicines may affect the way DOPTELET or DOPTELET Sprinkle works.

### What are the side effects of DOPTELET or DOPTELET Sprinkle?

DOPTELET or DOPTELET Sprinkle may cause serious side effects including **blood clots**. People with persistent or chronic ITP and people with certain blood clotting conditions may have an increased risk of developing blood clots.


Tell your healthcare provider right away if you get signs and symptoms of a blood clot, including:

- Swelling, pain, or tenderness in your legs
- Shortness of breath
- Chest pain
- Fast heartbeat
- Stomach (abdominal) pain or tenderness

**The most common side effects of DOPTELET or DOPTELET Sprinkle in children with persistent or chronic ITP are** viral infection, runny nose, cough, fever, pain in the mouth or throat.

These are not all of the possible side effects of DOPTELET or DOPTELET Sprinkle. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**Please see the DOPTELET or DOPTELET Sprinkle full [Prescribing Information](#)**



**Doptelet**<sup>®</sup>  
(avatrombopag)  
tablets

**Doptelet**<sup>®</sup>  
**Sprinkle**  
(avatrombopag) oral granules

## WHAT IS ITP?

Immune thrombocytopenia (ITP) is a blood disorder in which the immune system doesn't make enough platelets or attacks them by mistake.

ITP is categorized by how long the condition lasts: **newly diagnosed or acute** (up to 3 months), **persistent** (3 to 12 months), or **chronic** (more than 12 months).



### IMPORTANT SAFETY INFORMATION

#### What is DOPELET and DOPELET Sprinkle (avatrombopag)?

Doptelet is a prescription medicine used to treat low blood platelet counts in:

- children 1 year and older with persistent or chronic ITP when other treatments have not worked well enough

Please see Full Important Safety Information on pages 2-3 and the full [Prescribing Information](#).

## A few more things to know about ITP:

- 1 With too few platelets, the body can't prevent bleeding as it should.**  
With ITP, the immune system can attack platelets, leaving too few to protect against bleeding, which results in bruising.
- 2 Your child's body already tries to make more platelets.**  
It uses a natural signal called thrombopoietin (TPO). Doptelet and Doptelet Sprinkle mimic this signal to help raise platelet levels when other treatments haven't worked well enough.
- 3 There are different ways to treat ITP.**  
Some treatments focus on slowing platelet destruction (like steroids or IVIG), while others help the body make more platelets (like TPO-RA therapies).

Doptelet Sprinkle is indicated for patients aged  $\geq 1$  to  $<6$  years. Doptelet tablets are indicated for patients aged 6 and older. Doptelet Sprinkle oral granules that come in a bottle are not the same dosage form as Doptelet tablets that come in a blister card and cannot be directly substituted for each other.

IVIG=intravenous immunoglobulin; TPO-RA=thrombopoietin receptor agonist.

PLATELETS HELP STOP BLEEDING. IF THEY ARE LOW DUE TO PERSISTENT OR CHRONIC ITP, DOPELET AND DOPELET SPRINKLE MAY HELP BRING THEM UP!  
Flip the page to see how they help



### IMPORTANT SAFETY INFORMATION

DOPELET and DOPELET Sprinkle are different dosage forms of DOPELET.

DOPELET and DOPELET Sprinkle are not used to make platelet counts normal.

It is not known if DOPELET or DOPELET Sprinkle are safe and effective in children younger than 1 year of age.

**Doptelet**  
(avatrombopag)  
tablets

**Doptelet**  
**Sprinkle**  
(avatrombopag) oral granules

## DOPTELET AND DOPTELET SPRINKLE: FAST, LASTING, AND ONCE DAILY

Doptelet and Doptelet Sprinkle are TPO-RAs that treat low blood platelet counts in kids with persistent or chronic ITP. In a 3-month trial with children taking Doptelet or Doptelet Sprinkle and children who were not, Doptelet or Doptelet Sprinkle were shown to:

### LIFT PLATELETS

Raise platelet counts to **50,000 per microliter** or more for at least **2 consecutive weeks** of the 3-month trial. This was seen in 44 of 54 children taking Doptelet or Doptelet Sprinkle, compared with 0 of 21 taking placebo.

### ACT FAST

Lift platelet counts to 50,000 per microliter or more in as few as **8 days**. By Day 8, 30 of 54 children on Doptelet or Doptelet Sprinkle had platelet counts of 50,000 per microliter or more, compared with 0 of 21 people in the placebo group.

### AND LAST

In at least **6 out of the last 8 weeks of the study**, 15 of 54 children on Doptelet or Doptelet Sprinkle had a platelet count of 50,000 per microliter or more without the need for rescue treatment, compared with 0 of 21 taking placebo.



[CLICK HERE](#)  
for more about Doptelet  
and Doptelet Sprinkle.

Placebo group consists of individuals not receiving treatment.  
ITP=immune thrombocytopenia; TPO-RA=thrombopoietin receptor agonist.

### IMPORTANT SAFETY INFORMATION (continued)

Before you take **DOPTELET** or **DOPTELET Sprinkle**, tell your doctor about all your medical conditions, including if you:

- Have ever had a blood clot

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **DOPTELET** or **DOPTELET Sprinkle** may affect the way other medicines work, and other medicines may affect the way **DOPTELET** or **DOPTELET Sprinkle** works.

Please see Full Important Safety Information on pages 2-3 and the full [Prescribing Information](#).

## WHAT MAKES DOPTELET AND DOPTELET SPRINKLE DIFFERENT?

Doptelet and Doptelet Sprinkle are oral treatments that can give pediatric patients with persistent or chronic ITP a platelet count lift anytime, anywhere. Doptelet and Doptelet Sprinkle should be taken with food. Some food and liquid options for taking Doptelet Sprinkle are applesauce or orange juice. See full Instructions for Use for information on how to take Doptelet Sprinkle.




### No food-type restrictions

Doptelet tablet is a TPO-RA that must be taken with food and can be taken at any time. Because food won't impact the effectiveness of treatment, your child may have more flexibility with meal choices and schedules. The effectiveness of Doptelet Sprinkle also isn't impacted by food, but it must be taken with a soft food or liquid like applesauce or orange juice.



### No additional risk of liver damage

No additional liver function tests are required after starting Doptelet or Doptelet Sprinkle.




### No treatment injections or infusions

Doptelet and Doptelet Sprinkle are oral medications, so there are no extra doctor visits for treatment injections. (Platelet monitoring required.) Your child's doctor will adjust the dose of Doptelet or Doptelet Sprinkle based on platelet levels reached and will check your child's platelet counts weekly until their platelet counts reach  $\geq 50,000$  per microliter, then monthly after that, and for at least 4 weeks after stopping treatment.



### No eye exams

Doptelet and Doptelet Sprinkle don't require eye exams before starting or during treatment.



### Fits their schedule

Doptelet and Doptelet Sprinkle can be taken at any time, as long as they're taken at the same time every day. Remember to take the treatment as directed by your doctor. Tell your child's doctor about all the medicine your child takes, as their doctor may adjust their Doptelet or Doptelet Sprinkle dose.

### IMPORTANT SAFETY INFORMATION (continued)

#### What are the side effects of **DOPTELET** or **DOPTELET Sprinkle**?

**DOPTELET** or **DOPTELET Sprinkle** may cause serious side effects including **blood clots**. People with persistent or chronic ITP and people with certain blood clotting conditions may have an increased risk of developing blood clots.

**Doptelet**  
(avatrombopag)  
tablets

**Doptelet**  
**Sprinkle**  
(avatrombopag) oral granules

## DOSING THAT CAN WORK WITH YOUR CHILD'S SCHEDULE

Doptelet and Doptelet Sprinkle can help match age-based needs in children.

### DOPTELET

**A 20-mg tablet taken with food.** For patients aged **≥6 years**.

### DOPTELET SPRINKLE

Coming soon **as a 10-mg capsule** containing oral granules for mixing with a soft food or liquid.\*

Click the link in the blue box to see the instructions for use. For patients **aged ≥1 to <6 years**.

#### \*EASY-TO-TAKE DOSING FOR CHILDREN

You can mix Doptelet Sprinkle with soft foods like applesauce or yogurt, or with drinks like water, whole milk, or orange juice. Choose what works best for your child.

DOPTELET may be affected by other drugs and may require a dose adjustment when coadministered with other drugs. Tell your child's doctor about other prescription or nonprescription medications or dietary supplements. If your child is taking CYP2C9 and CYP3A4 inhibitors such as certain antibiotics, antifungals, or antidepressants, their doctor will start them on 20 mg of Doptelet tablet or 10 mg of Doptelet Sprinkle 3 times a week. If they are taking CYP2C9 and CYP3A4 inducers such as antibiotics, antifungals, or antiepileptic drugs, they will start on 40 mg of Doptelet tablet or 20 mg of Doptelet Sprinkle once a day.

[CLICK HERE](#)  
to learn more about dosing for kids.

DOPTELET TABLETS CAN BE  
TAKEN WITH ANY FOOD,  
EVEN PIZZA—YUM!

Food required.



## IMPORTANT SAFETY INFORMATION FOR DOPTELET AND DOPTELET SPRINKLE

### Serious side effects

Doptelet and Doptelet Sprinkle are thrombopoietin (TPO) receptor agonists and TPO receptor agonists have been associated with thrombotic and thromboembolic complications in patients with chronic liver disease or immune thrombocytopenia. Two patients experienced serious adverse reactions to Doptelet or Doptelet Sprinkle during the core phase: thrombocytosis and headache.

- Swelling, pain, or tenderness in their legs
- Shortness of breath
- Chest pain
- Fast heartbeat
- Stomach (abdominal) pain or tenderness

### Common side effects

The most common side effects of Doptelet or Doptelet Sprinkle when used to treat low blood platelet counts in children 1 year and older with persistent or chronic immune thrombocytopenia (ITP) are: viral infection (including viral upper respiratory infection, viral infection, COVID-19, parainfluenza virus infection, and rhinovirus infection), runny nose, cough, fever, pain in the mouth or throat.

**These are not all the possible side effects of Doptelet and Doptelet Sprinkle. Talk to your child's doctor if any side effect bothers your child or does not go away.**

#### IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider right away if you get signs and symptoms of a blood clot, including:

- Swelling, pain, or tenderness in your legs
- Shortness of breath
- Chest pain
- Fast heartbeat
- Stomach (abdominal) pain or tenderness

Please see Full Important Safety Information on pages 2-3 and the full [Prescribing Information](#).

#### IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of **DOPTELET** or **DOPTELET Sprinkle** in children with persistent or chronic ITP are viral infection, runny nose, cough, fever, pain in the mouth or throat.

**Doptelet**<sup>®</sup>  
(avatrombopag)  
tablets

**Doptelet**<sup>®</sup>  
**Sprinkle**  
(avatrombopag) oral granules



## GET STARTED WITH DOPTELET CONNECT

Doptelet Connect is designed as a single point of contact to help you understand your child's insurance coverage after your child's healthcare provider prescribes Doptelet or Doptelet Sprinkle. Doptelet Connect can also answer questions about coverage and financial assistance options.

**After your doctor prescribes Doptelet or Doptelet Sprinkle, Doptelet Connect can provide you with information on patient assistance options that may be available to help you access Doptelet or Doptelet Sprinkle. These offerings include:**



Evaluating your child's insurance coverage



Accessing the Patient Assistance Program\*  
\*For eligible patients.



Exploring financial assistance options



Assessing the Doptelet Free Trial Offer

### DO YOU WANT TO LEARN MORE ABOUT PATIENT SUPPORT PROGRAMS?

For additional assistance, call Doptelet Connect at **1-833-368-2663**, Monday–Friday, 8:30 AM–7 PM ET.

### CLICK HERE

for more information on financial assistance.

### IMPORTANT SAFETY INFORMATION (continued)

These are not all of the possible side effects of DOPTELET or DOPTELET Sprinkle. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see the DOPTELET or DOPTELET Sprinkle full Prescribing Information.

Please see Full Important Safety Information on pages 2-3 and the full [Prescribing Information](#).

## WHAT TO EXPECT ONCE YOUR CHILD HAS BEEN PRESCRIBED DOPTELET OR DOPTELET SPRINKLE

Your child's prescription is sent to a pharmacy and insurance coverage is confirmed



Your child's prescription will be filled at either a specialty or clinic pharmacy—based on your insurance provider—and you'll be informed about any out-of-pocket costs.

The pharmacy will call you



It's important to answer the pharmacy's call to confirm your child's prescription details. If you miss the call, call them back as soon as possible. Doptelet or Doptelet Sprinkle can't be delivered until you've spoken to a pharmacist to confirm the order.

Your doctor and pharmacy will set up future refills



Stay in touch with your child's doctor and pharmacy to manage future refills and avoid any interruptions in treatment.  
**Note: The pharmacy will call you to confirm each delivery.**



### BE SURE TO ANSWER THE CALL

Your pharmacy may call from an 800 number—be sure to answer to avoid delaying the delivery of your child's medicine.

**Doptelet**<sup>®</sup>  
(avatrombopag)  
tablets

**Doptelet**<sup>®</sup>  
**Sprinkle**  
(avatrombopag) oral granules

## GETTING READY TO TALK TO YOUR CHILD'S DOCTOR?

Treating ITP is a team effort. Setting clear goals early can help create a plan that fits your child's needs and your family's routine.

### Why treatment goals matter:

- Reaching your target platelet counts can help support treatment goals in ITP
- Your child's doctor will adjust the dose of Doptelet or Doptelet Sprinkle based on platelet levels reached and will check your child's platelet counts weekly until their platelet counts reach 50,000 per microliter, then monthly after that, and for at least 4 weeks after stopping treatment
- Once goals are met, blood tests may happen less often. Talk to your child's doctor for more information

### The goal? Target platelet counts.

Raising platelets to the target level of 50,000 per microliter or more may help protect against bleeding

### Questions to consider:

- What is your child's most important treatment goal and why?
- Could this treatment affect your child's eating habits, including the types of food and what time they eat?
- What type of treatment do you think might work best for your child?
- Do you need help with access or financial support?

**TIP**  
Use the back page to write down your questions and answers.



Please see Full Important Safety Information on pages 2-3 and the full [Prescribing Information](#).

## ITP EXPLAINED IN A KID-FRIENDLY WAY

### What is ITP?

ITP is short for immune thrombocytopenia (try saying that five times fast), and it occurs when your body has fewer platelets than it needs. ITP is categorized by how long the condition lasts: newly diagnosed or acute (up to 3 months), persistent (3 to 12 months), or chronic (more than 12 months).

### What might happen if you have ITP?

- You might feel more tired than usual
- You might get bruises easily
- You could see red or purple spots on your skin
- You might need to take breaks or skip some activities

These are all normal parts of having ITP. Doptelet and Doptelet Sprinkle do not treat these signs and symptoms.

### How to talk about ITP with others

Sometimes people might not know what ITP is.

But you can help them understand by saying things like:

"I HAVE SOMETHING CALLED ITP. IT MAKES ME GET BRUISES MORE EASILY."

"SOMETIMES I NEED TO REST TO TAKE CARE OF MYSELF."

"MY DOCTOR HELPS ME KEEP AN EYE ON IT."



**Doptelet**  
(avatrombopag)  
tablets

**Doptelet**  
**Sprinkle**  
(avatrombopag) oral granules

# NOTES FOR YOUR CHILD'S DOCTOR VISIT

Use this page to write down your thoughts and questions on treatment goals to discuss with your child's doctor.

## What matters most to my child:

---

---

## Questions or concerns I want to talk about:

---

---

### **SIGN UP TO LEARN MORE**

Receive tools and tips designed to help support your family.



Doptelet is a registered trademark of AkaRx, Inc., a Sobi company.  
Sobi is a registered trademark of Swedish Orphan Biovitrum AB (publ)  
Doptelet Connect is a trademark of AkaRx, Inc.  
© 2025 Sobi, Inc. – All rights reserved.  
PP-29112 08/25

**Doptelet**<sup>®</sup>  
(avatrombopag)  
tablets

**Doptelet**<sup>®</sup>  
**Sprinkle**  
(avatrombopag) oral granules