

HOW TO TAKE DOPTELET

Your guide to taking DOPTELET before your procedure



What is DOPTELET (avatrombopag)?

DOPTELET is a prescription medicine used to treat low blood platelet counts in adults with long-lasting (chronic) liver disease who are scheduled to have a medical or dental procedure.

DOPTELET is not used to make platelet counts normal in adults with chronic liver disease. It is not known if DOPTELET is safe and effective in children.

SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTELET in people with chronic liver disease are fever, stomach (abdominal) pain, nausea, headache, tiredness, and swelling of the hands or feet. These are not all the possible side effects of DOPTELET.

DOPTELET is not used to make platelet counts normal in adults with chronic liver disease or chronic immune thrombocytopenia. It is not known if DOPTELET is safe and effective in children.

Please see Important Safety Information on back page and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at DOPTELET.com.

Doptelet[®]
(avatrombopag) tablets

How should I take DOPTELET?

You should take DOPTELET® (avatrombopag) just as your healthcare provider prescribed.

Three things you should know about taking DOPTELET:



ONCE A DAY. Take your dose of DOPTELET once a day.



WITH FOOD. Take DOPTELET every day with food.

**5
DAYS**

FOR 5 DAYS. Take DOPTELET for 5 days in a row as directed by your healthcare provider.

- If you are taking DOPTELET prior to a scheduled medical procedure and you miss a dose, contact your healthcare provider for further dosing instructions.
- Complete all 5 days of medication.

Starting on the right day is important.

Someone from your healthcare team, most likely the specialty pharmacy, will contact you to remind you when to start taking DOPTELET.

If you have any questions about how or when to take DOPTELET, call your healthcare provider. You can also call your specialty pharmacy, or Dova 1Source at 1-833-DOVA-ONE (1-833-368-2663) and they will help you.

SELECTED IMPORTANT SAFETY INFORMATION

The most common side effects of DOPTELET are fever, stomach (abdominal) pain, nausea, headache, tiredness, swelling of the hands or feet. These are not all the possible side effects of DOPTELET.

Please see Important Safety Information on back page and accompanying Full Prescribing Information for DOPTELET (avatrombopag), including Patient Information; also available at DOPTELET.com.

DOPTELET 40 mg—2 tablets a day for 5 days



DOPTELET 60 mg—3 tablets a day for 5 days



Your DOPTELET schedule will look like this.

Speak to your healthcare provider about the exact day you should start DOPTELET. Your specialty pharmacy will contact you to remind you of those days. You can use the schedule below to write in your days. This may help you remember.

DAY 1: _____ DAY 4: _____

DAY 2: _____ DAY 5: _____

DAY 3: _____ PROCEDURE DAY: _____



IMPORTANT SAFETY INFORMATION FOR DOPTELET

What are the possible side effects of DOPTELET?

- DOPTELET may cause serious side effects, including blood clots. People with chronic liver disease and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you have signs and symptoms of a blood clot, including swelling, pain or tenderness in your leg; shortness of breath; chest pain; fast heartbeat; or stomach (abdominal) pain or tenderness.
- The most common side effects of DOPTELET are fever, stomach (abdominal) pain, nausea, headache, tiredness, swelling of the hands or feet. These are not all the possible side effects of DOPTELET.

Before you take DOPTELET, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTELET may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTELET.
- are breastfeeding or plan to breastfeed. It is not known if DOPTELET passes into your breast milk. Do not breastfeed during your treatment with DOPTELET and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTELET may affect the way other medicines work, and other medicines may affect the way DOPTELET works.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For product information and adverse event reports, please contact:

Dova Medical Information Call Center

Phone: 1-844-506-3682

E-mail: medinfo@Dova.com

Reference: DOPTELET [package insert]. Durham, NC: Dova Pharmaceuticals, Inc; 2019.

Please see accompanying Full Prescribing Information, including Patient Information; also available at Doptelet.com.

If you have any questions about how or when to take DOPTELET, call your healthcare provider. You can also call your specialty pharmacy, or Dova 1Source at 1-833-DOVA-ONE (1-833-368-2663) and they will help you.



DOPTELET is a registered trademark of AkaRx, Inc.

©2019 Dova Pharmaceuticals, Inc.

All rights reserved. PM-US-DOP-0157 Printed in USA.

Doptelet
(*avatrombopag*) tablets